

xarvio SCOUTING Accuracy Report

5/20/2019



| Lokaler Name | Wissenschaftlicher Name | Gesamt | Keimblatt- stadium | Frühes Stadium | Ausgereift | Verzweigung | Blüte |
|-------------------------------|---------------------------|--------|-----------------------|-------------------|------------|-------------|-------|
| Amarant; Dorniger | Amaranthus spinosus | 96% | 83% | 97% | 75% | 96% | N/A |
| Amarant; Fuchsschwanz- | Amaranthus palmeri | 85% | 72% | 88% | 100% | N/A | N/A |
| Amarant; Grünlicher | Amaranthus lividus | 96% | 100% | 96% | 87% | N/A | N/A |
| Amarant; Liegender | Amaranthus viridis | 97% | 96% | 96% | 98% | 99% | N/A |
| Amarant; Niederliegender | Amaranthus blitoides | 87% | 50% | 86% | 92% | 89% | N/A |
| Amarant; Zurückgekrümmter | Amaranthus retroflexus | 87% | 66% | 91% | 89% | 73% | 67% |
| Ambrosie; Beifussblättrige | Ambrosia artemisiifolia | 90% | 88% | 89% | 93% | 50% | N/A |
| Ampfer; Krauser | Rumex crispus | 61% | 56% | 69% | 50% | N/A | N/A |
| Ampfer; Sauer- | Rumex acetosa | 94% | 100% | 94% | 95% | N/A | N/A |
| Archangel | Eupatorium odoratum | 84% | 78% | N/A | 100% | N/A | N/A |
| Asthma plant | Chamaesyce hirta | 97% | 87% | 98% | 98% | 99% | 99% |
| Berufskraut; Argentinisches | Erigeron bonariensis | 98% | 70% | 100% | 98% | 100% | N/A |
| Berufskraut; Kanadisches | Erigeron canadensis | 83% | 67% | 88% | 65% | N/A | 100% |
| Berufskraut; Weißliches | Erigeron sumatrensis | 100% | N/A | 100% | 100% | 100% | N/A |
| Besenradmelde | Bassia scoparia | 89% | 100% | 86% | 92% | 100% | N/A |
| Bingelkraut; Einjähriges | Mercurialis annua | 93% | 94% | 92% | 96% | N/A | 100% |
| Black pigweed | Trianthema portulacastrum | 90% | 79% | 81% | N/A | N/A | 83% |
| Blattblume | Phyllanthus tenellus | 98% | N/A | 100% | 100% | 100% | N/A |
| Brazil pusley | Richardia brasiliensis | 94% | 90% | 98% | 79% | 67% | 75% |
| Brennnessel; Kleine | Urtica urens | 87% | 73% | 88% | 88% | N/A | N/A |
| Buttonweed; Broadleaf | Borreria latifolia | 100% | 100% | N/A | 92% | 50% | N/A |
| Chocolateweed | Melochia corchorifolia | 87% | 67% | 100% | 100% | N/A | N/A |
| Cirsium segetum | Cirsium segetum | 93% | 94% | 87% | 100% | N/A | N/A |
| Commeline; Bengalische | Commelina benghalensis | 98% | 97% | 96% | 97% | 97% | N/A |
| Dreibiß; Niederliegender | Tridax procumbens | 97% | 97% | 99% | 96% | 93% | 100% |
| Ehrenpreis; Glänzender | Veronica polita | 97% | 100% | 95% | 100% | N/A | N/A |
| Ehrenpreis; Persischer | Veronica persica | 86% | 63% | 90% | 86% | 96% | 50% |
| Erdrauch; Gemeiner | Fumaria officinalis | 82% | 71% | 83% | 86% | N/A | 100% |
| Franzosenkraut; Behaartes | Galinsoga quadriradiata | 96% | 94% | 96% | 96% | N/A | N/A |
| Franzosenkraut; Kleinblütiges | Galinsoga parviflora | 97% | 100% | 92% | 96% | N/A | 100% |
| Gänsedistel; Acker- | Sonchus arvensis | 92% | 89% | 94% | 90% | N/A | N/A |

xarvio SCOUTING Accuracy Report

5/20/2019



| Lokaler Name | Wissenschaftlicher Name | Gesamt | Keimblatt- stadium | Frühes Stadium | Ausgereift | Verzweigung | Blüte |
|------------------------------|---------------------------|--------|-----------------------|-------------------|------------|-------------|-------|
| Gänsedistel; Dornige | Sonchus asper | 76% | 64% | 85% | 72% | N/A | N/A |
| Gänsedistel; Kohl- | Sonchus oleraceus | 80% | 88% | 73% | 85% | 100% | N/A |
| Gänsefuß; Vielsamiger | Lipandra polysperma | 83% | N/A | 91% | 83% | N/A | N/A |
| Gänsefuß; Weißer | Chenopodium album | 95% | 88% | 95% | 98% | 80% | 78% |
| Gauchheil; Roter | Anagallis arvensis | 79% | 77% | 82% | 78% | 100% | N/A |
| Giftbeere | Nicandra physalodes | 98% | 100% | 97% | 99% | 100% | 100% |
| Gooseweed | Sphenoclea zeylanica | 96% | 81% | 98% | 99% | 100% | N/A |
| Hairy indigo | Indigofera hirsuta | 100% | 99% | 100% | 100% | 100% | N/A |
| Hederich | Raphanus raphanistrum | 98% | 99% | 98% | 99% | 100% | 98% |
| Hirtentäschelkraut; Gemeines | Capsella bursa-pastoris | 90% | 75% | 90% | 90% | 100% | 50% |
| Hispid starbur | Acanthospermum hispidum | 99% | 100% | 100% | 97% | N/A | N/A |
| Hornklee; Gewöhnlicher | Lotus corniculatus | 90% | 85% | 94% | 94% | 100% | N/A |
| Hornkraut; Knäuel- | Cerastium glomeratum | 93% | 67% | 95% | 100% | N/A | N/A |
| Hundskamille; Acker- | Anthemis arvensis | 75% | 77% | 82% | 63% | 100% | N/A |
| Hundspetersilie; Gemeine | Aethusa cynapium | 91% | 100% | 93% | 89% | N/A | N/A |
| Ipomoea grandifolia | Jacquemontia grandifolia | 100% | 100% | 100% | 100% | 86% | N/A |
| Johanniskraut; Echtes | Hypericum perforatum | 98% | N/A | 95% | 100% | 100% | N/A |
| Kamille; Duftlose | Tripleurospermum inodorum | 74% | 75% | 72% | 75% | 80% | N/A |
| Kamille; Echte | Matricaria chamomilla | 91% | 93% | 92% | 90% | 100% | N/A |
| Klettenlabkraut | Galium aparine | 90% | 88% | 90% | 89% | 93% | 100% |
| Knöterich; Ampfer- | Persicaria lapathifolia | 82% | 75% | 77% | 88% | 100% | N/A |
| Knöterich; Floh- | Persicaria maculosa | 71% | 75% | 67% | 77% | 100% | N/A |
| Knöterich; Vogel- | Polygonum aviculare | 82% | 47% | 88% | 85% | 79% | N/A |
| Knöterich; Winden- | Fallopia convolvulus | 92% | 88% | 94% | 94% | 89% | 100% |
| Kornblume | Cyanus segetum | 92% | 87% | 94% | 88% | 100% | N/A |
| Kreuzkraut; Gemeines | Senecio vulgaris | 89% | 92% | 88% | 91% | 100% | 86% |
| Lady's thumb; Oriental | Persicaria longisetata | 100% | 100% | 100% | N/A | N/A | N/A |

xarvio SCOUTING Accuracy Report

5/20/2019



| Lokaler Name | Wissenschaftlicher Name | Gesamt | Keimblatt- stadium | Frühes Stadium | Ausgereift | Verzweigung | Blüte |
|-------------------------------|-------------------------|--------|-----------------------|-------------------|------------|-------------|-------|
| Lederbalsam | Ageratum conyzoides | 100% | N/A | 100% | N/A | N/A | N/A |
| Lein; Gemeiner | Linum usitatissimum | 100% | 100% | 100% | 100% | N/A | N/A |
| Löwenzahn; Gemeiner | Taraxacum officinale | 90% | 91% | 88% | 92% | 100% | 100% |
| Malve; Kleinbütiige | Malva pusilla | 90% | 67% | 100% | 92% | 100% | N/A |
| Malve; Weg- | Malva neglecta | 82% | N/A | 75% | 100% | 100% | N/A |
| Mehlblume | Eclipta prostrata | 95% | 100% | 97% | 83% | N/A | N/A |
| Mohn; Klatsch- | Papaver rhoeas | 86% | 83% | 89% | 81% | N/A | N/A |
| Möhre; Wilde | Daucus carota | 88% | 86% | 91% | 80% | N/A | N/A |
| Nachtschatten; Amerikanischer | Solanum americanum | 98% | N/A | N/A | 98% | N/A | N/A |
| Nachtschatten; Schwarzer | Solanum nigrum | 89% | 78% | 94% | 86% | 100% | 67% |
| Papageienblatt; Sitzendes | Alternanthera sessilis | 69% | N/A | 68% | 61% | N/A | N/A |
| Portulak | Portulaca oleracea | 94% | 98% | 88% | 98% | 85% | 99% |
| Primrose; Water | Ludwigia adscendens | 53% | N/A | 64% | 58% | N/A | 100% |
| Primrose-willow; Mexican | Ludwigia octovalvis | 91% | 83% | 90% | 82% | 99% | N/A |
| Prunkwinde; Purpur- | Ipomoea purpurea | 90% | 50% | 100% | 83% | N/A | N/A |
| Raps | Brassica napus (winter) | 92% | 93% | 93% | 88% | 100% | 100% |
| Samtpappel | Abutilon theophrasti | 95% | 93% | 98% | 89% | N/A | 100% |
| Schachtelhalm; Acker- | Equisetum arvense | 80% | 84% | 80% | 62% | 80% | N/A |

xarvio SCOUTING Accuracy Report

5/20/2019



| Lokaler Name | Wissenschaftlicher Name | Gesamt | Keimblatt- stadium | Frühes Stadium | Ausgereift | Verzweigung | Blüte |
|-----------------------------------|-------------------------|--------|-----------------------|-------------------|------------|-------------|-------|
| Schafgarben (Gattung) | Achillea | 90% | 86% | 87% | 100% | N/A | N/A |
| Sicklepod | Senna obtusifolia | 99% | 100% | 100% | 99% | 100% | 67% |
| Sidafaserpflanze | Sida rhombifolia | 88% | 71% | 96% | 88% | N/A | N/A |
| Smartweed; Prickly | Polygonum bungeanum | 98% | 88% | 100% | 99% | N/A | N/A |
| Sonnenblume | Helianthus annuus | 84% | 75% | 93% | 60% | N/A | N/A |
| Spitzklette; Gewöhnliche | Xanthium strumarium | 90% | 91% | 94% | 90% | 67% | N/A |
| Stechapfel; Gemeiner | Datura stramonium | 88% | 70% | 93% | 91% | N/A | 100% |
| Stiefmütterchen; Acker- | Viola arvensis | 85% | 80% | 88% | 85% | 100% | 45% |
| Stinkkassie | Senna occidentalis | 100% | 100% | 100% | 99% | 100% | 100% |
| Storchschnabel; Schlitzblättriger | Geranium dissectum | 92% | 90% | 93% | 91% | 100% | N/A |
| Sumpflieb | Limnocharis flava | 66% | 66% | 63% | N/A | N/A | N/A |
| Tagblume; Asiatische | Murdannia keisak | 98% | 97% | 99% | 100% | N/A | N/A |
| Tagblume; Gewöhnliche | Commelina communis | 95% | 100% | 98% | 69% | N/A | N/A |
| Taubnessel; Rote | Lamium purpureum | 78% | 77% | 77% | 78% | 100% | 86% |
| Taubnessel; Stängelumfassende | Lamium amplexicaule | 88% | 63% | 87% | 90% | 95% | N/A |
| Traubenkraut; Dreilappiges | Ambrosia trifida | 93% | 94% | 85% | 100% | N/A | N/A |
| Trichterwinde; Efeublättrige | Ipomoea hederifolia | 97% | 92% | N/A | 100% | 100% | N/A |
| Trichterwinde; Gefiederte | Ipomoea quamoclit | 100% | 100% | N/A | 100% | 100% | N/A |
| Vergissmeinnicht; Acker- | Myosotis arvensis | 92% | 90% | 92% | 94% | N/A | N/A |
| Vogelmiere | Stellaria media | 93% | 89% | 85% | 99% | 98% | 100% |
| Waterhemp; Common | Amaranthus tamariscinus | 74% | 100% | 69% | 79% | N/A | N/A |
| Wegerich; Spitz- | Plantago lanceolata | 76% | N/A | 81% | 73% | N/A | N/A |
| Wicke; Futter- | Vicia sativa | 78% | N/A | 89% | 89% | N/A | N/A |
| Winde; Acker- | Convolvulus arvensis | 84% | 81% | 87% | 83% | 100% | N/A |
| Wolfsmilch; Bechertragende | Euphorbia heterophylla | 85% | 96% | 72% | 86% | 95% | N/A |
| Wunderbaum | Ricinus communis | 99% | N/A | 99% | 99% | N/A | N/A |
| Zweizahn; Behaarter | Bidens pilosa | 98% | 95% | 100% | 95% | 93% | N/A |
| Zweizahn; Schwarzfrüchtiger | Bidens frondosa | 92% | 75% | 93% | 100% | N/A | N/A |